

SIMULTANEOUS SCREENING AND IDENTIFICATION OF SEQUENCE ALTERATIONS FROM AMPLIFIED TARGET

5

FIELD OF THE INVENTION

The invention relates to methods for screening for and identifying sequence alterations in nucleic acids.

10

BACKGROUND OF THE INVENTION

15

The ability to scan and identify sequence alterations has widespread applications in many areas including genetics, immunology, infectious disease, oncology, epidemiology and forensics. Mutations leading to cancer can arise in a number of different genes and in different positions within the same gene. Gene variants also can be the source of hereditary diseases. Scanning and identification of such nucleic acid alterations have important implications for diagnosis and prognosis and for guiding therapy.

20

Several methods exist for scanning for the presence or absence of sequence variants. Denaturing Gradient Gel electrophoresis (DGGE) takes advantage of differences in the melting location of slightly different nucleotide sequences on a gradient gel during electrophoresis. (Mers et al., *Methods Enzymol.* 155:501-527 (1987); Abrams et al., *Genomics* 7:463-475 (1990); Vijg et al., U.S. Patent No. 5,814,491) The process is generally performed on fragments of a nucleic acid of interest produced through polymerase chain reaction (PCR). Similarly, differences in mobility during electrophoresis of single strands of nucleic acid is used in the technique known as Single-Strand Conformation Polymorphism (SSCP). (Orita et al., *Genomics* 5:874-879 (1989)) This method takes advantage of different conformations assumed by nucleic acids having slightly different sequences under non-denaturing conditions. But, while these methods can identify nucleic acid sequences with alterations, the nature and exact location of the alteration(s) must be subsequently determined by other techniques.

30

35

Identification of the location and nature of an alteration in a nucleic acid sequence may be determined by direct sequencing. However, this process is generally very labor intensive and time-consuming. Other means of detecting known sequence alterations involve the use of oligonucleotides which hybridize to the specific altered sequence. (Conner et al., *Proc. Nat. Acad. Sci.*, 80:278-282 (1983)) This method becomes untenable, however, when an

alteration is unknown. Other methods utilizing restriction enzymes have been developed to identify alterations in a sequence without directly sequencing the nucleic acid. Unfortunately, these enzymes recognize only a limited number of restriction sequences. And none of these methods provides for quick determination of which alteration is actually present.

It is apparent that there is a need for a fast and simple method of screening nucleic acids for alterations and determining the precise location and nature of such an alteration.

Accordingly, it is an object of the present invention to provide novel methods of screening for sequence alterations in a target nucleic acid. The invention provides methods utilizing probes which hybridize to nucleic acid to detect alterations in a target sequence as compared to a control.

It is a further object of the invention to provide methods of screening for and determining the location of a sequence alteration in a target nucleic acid in a single step. These methods utilize probes which hybridize to nucleic acid to detect the existence and location of alterations in a target sequence.

An additional object of the invention is to provide methods of screening, determining the location and determining the nature of a sequence alteration in a target nucleic acid in a single step. These methods provide for the identification of specific base changes in the target sequence.

SUMMARY OF THE INVENTION

In accordance with the objectives outlined above, the present invention provides methods of screening and identifying alterations in a target nucleic acid sequence as compared with a control nucleic acid. Probes are produced which are complementary to and, therefore, hybridize to overlapping regions of a control nucleic acid. The methods are based on the fact that probes directed to a control sequence denature from a target sequence at a different temperature than from the control when the target sequence has an alteration (such as a point mutation) as compared with the control at the location at which the probe hybridizes.

In one aspect of the present invention, a method is provided for identifying a sequence alteration in a target nucleic acid as compared to a control nucleic acid. The method entails hybridizing a plurality of nucleic acid probes with said target nucleic acid, wherein said probes are complementary to different overlapping regions of said control nucleic acid. The melting temperature (T_m) of at least two overlapping probes is determined, as well as the difference

between the melting temperature of each probe from the target nucleic acid and the control nucleic acid (ΔT_m). The difference between the ΔT_m s of overlapping probes is determined as an indication of whether or not a sequence alteration exists in the target nucleic acid as compared to the control nucleic acid. In one embodiment, as few as two probes are required to examine a specific sequence alteration in target nucleic acid as compared with control nucleic acid.

In the method described above, the difference in ΔT_m between at least two overlapping probes indicates the location of a nucleotide difference in the target nucleic acid as compared to the control nucleic acid. In addition, the difference in ΔT_m between at least two overlapping probes indicates a substitution in the target nucleic acid sequence as compared to the control nucleic acid. Furthermore, the difference in ΔT_m between at least two overlapping probes indicates the type of nucleotide substituted in the target nucleic acid sequence as compared to the control nucleic acid.

In another aspect of the invention, provided herein is a method for identifying a sequence alteration in a target nucleic acid as compared to a control nucleic acid. The method involves hybridizing a plurality of nucleic acid probes with the target nucleic acid; a first set of probes is complementary to regions of the control nucleic acid separated by one or more nucleotides and at least a second set of probes is complementary to regions of the control separated by one or more nucleotides. The regions complementary to the second set of probes include the nucleic acids separating the first set of probes and are overlapping with the regions complementary to the first set of probes. The method further entails determining the melting temperature (T_m) of at least two overlapping probes from the target nucleic acid, determining for these at least two overlapping probes the difference between the T_m from the target nucleic acid and the T_m from the control nucleic acid (ΔT_m), and determining the difference in determined ΔT_m between overlapping probes. The difference ΔT_m between overlapping probes provides an indication of the presence or absence of a sequence alteration in the target nucleic acid as compared to the control nucleic acid. Preferably only two sets of probes are used.

Further to the method just described, the difference in ΔT_m between at least two overlapping probes indicates the location of a nucleotide difference in the target nucleic acid as compared to the control nucleic acid. In addition, the difference in ΔT_m between at least two overlapping probes indicates a substitution in the target nucleic acid sequence as compared to the control nucleic acid. Furthermore, the difference in ΔT_m between at least two overlapping probes indicates the type of nucleotide substituted in the target nucleic acid sequence as compared to the control nucleic acid.

Also with respect to the method just described, in one embodiment only three probes are used, two from the first set which are complementary to adjacent regions and one from the second probe set which overlaps each of the probes from the first probe set. In this latest case, when the probes from the first probe set have a ΔT_m of zero and the probe from the second probe set has a ΔT_m other than zero, an alteration in the target is indicated as being in the nucleotides of the control nucleic acid sequence separating the regions to which the probes from the first probe set are complementary. In the case that one nucleotide separates adjacent regions to which the probes of the first probe set are complementary, a ΔT_m of zero for two adjacent (non-overlapping) probes and a ΔT_m other than zero for a probe which overlaps both adjacent probes indicated that the alteration is at the nucleotide separating the adjacent regions. In this last case, the ΔT_m of the overlapping probe indicates both a substitution as the sequence alteration and the type of substitution.

DESCRIPTION OF THE DRAWINGS

Figure 1 depicts schematically how overlapping probes are walked along a complementary single-stranded target such that each probe shares a common sequence with a probe placed before and after it. This figure also shows actual experimental data illustrating the relative destabilization of overlapping probes in response to single nucleotide alterations in the target at different locations in the overlapping region.

Figure 2 depicts an alternative embodiment of the present invention in which a set of probes are walked along the complementary target, separated by a single base between flanking probes, and a second set of probes is similarly walked along the target, separated by a single base and centered on the bases separating the first set of probes. This figure also presents predicted changes in melting temperature of adjacent and overlapping probes in response to a single base alteration in the target at different locations.

Figure 3 shows the effects on the melting temperature of a 15mer oligonucleotide probe from an otherwise complementary target in response to different nucleotide pairings at the center position of the probe. A Unique melting temperature is seen for each pairing.

Figure 4 shows the effects on the melting temperature of a 20mer probe from an otherwise complementary target in response to a single G:T mismatch at different locations along the length of the probe. A unique melting temperature is seen for each location.

Figure 5 shows ΔT_m for G:T mismatches at 20 different positions within the hybridization region of a 20mer probe to a target. This figure illustrates the increasing destabilization of a

duplex as a G:T mismatch is moved from the end position of the probe to the center. At each position, T_m was determined for a completely complementary duplex and the same duplex with a G:T mismatch.

Figure 6 shows how the position of a mismatch can be localized using the ΔT_m 's of 2 overlapping probes to determine ΔT_m . The two probes had an overlapping hybridization region of 10 nucleotides on the targets. When the mismatch is positioned at a similar distance from the end of each probe, the ΔT_m is at or near zero. In comparison, if the mismatch is positioned near the end of one probe and the center of another, the ΔT_m is at its greatest. The direction of the ΔT_m sign (i.e., + or -) indicates at which end of the overlapping region the mismatch is positioned.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides methods of screening for and identifying sequence alterations in a target nucleic acid. These methods utilize probes which are complementary to a control nucleic acid. The temperature of thermal denaturation (melting) of these probes when hybridized with the target is compared with the melting temperatures of the same probes from the control. Surprisingly, determination of melting temperature of the individual probes from the target and the control provide sufficient information to determine not only the location of an alteration, but the nature of a nucleotide substitution. A difference in melting temperature of a probe detects an alteration and provides information about the nature of the alteration. Differences in melting temperatures of overlapping probes provides information to localize the alteration. Advantageously, the screening and identification of the sequence alteration can be obtained in a single thermal denaturation step.

It is contemplated that the present invention may be applied to virtually any nucleic acid for which the sequence is known. The methods are particularly useful for identifying single nucleotide substitutions in a target.

By "melting temperature" (T_m) is meant the temperature at which a probe dissociates from a nucleic acid to which it is hybridized under defined conditions. The conditions are preferably chosen so as to provide the maximum change in T_m if a mismatch is present. Examples of preferred hybridization conditions are conditions used for asymmetric amplification by PCR, so that no sample handling is necessary. The same solution can then be amplified and melted in the same instrument to determine T_m values. An example of asymmetric amplification and melting analysis is found in Lay and Wittwer, *Clin. Chem.* 43:2262-2267 (1997), which is incorporated herein in its entirety. In one embodiment of the present invention, asymmetric amplification is used to produce an excess of one strand of a target or control, so that the

overproduced product hybridizes to the probes rather than to its complementary product strand.

5 In another embodiment of the invention, single strand production and melting analysis is performed as separate steps. Single strand nucleic acid may be produced by asymmetric PCR or other techniques known in the art. Examples of such techniques include self sustained sequence replication (3SR) and strand displacement amplification (SDA). The single strand target or control is then combined with the probes and T_m values are determined. In this embodiment, any probe hybridization solution may be used, such solutions being well known in the art.

10 By "probe" is meant any sequence of molecules which is complementary to the control. By "complementary" is meant that the probe has a specific sequence of molecules which specifically hybridizes on a residue-by-residue basis with a sequence of nucleic acid. Such molecules include nucleic acids like DNA or RNA, and probes can contain both deoxy- and ribonucleic acids (oligonucleotides). The probes may have a ribose-phosphate backbone typical of naturally occurring DNA or RNA, but may also contain modifications in such a ribose-phosphate backbone. Probes may also be in other forms, for example as peptide nucleic acids.

15 Means for producing probes having a desired sequence are well known in the art. Preferably, each probe has the same number of nucleotides. Preferably, each probe is between 10 and 50 nucleic acids in length, more preferably between 10 and 30 nucleic acids, still more preferably, from 15 to 20 nucleic acids. In one embodiment, each probe is a 15-base nucleic acid. In another embodiment, each probe is a 20-base nucleic acid.

20 In one embodiment of the invention, the probes are completely overlapping. By "overlapping" is meant that at least two probes hybridize with the same one or more nucleotides of the control nucleic acid. This embodiment will be referred to as the complete overlapping embodiment. In a preferred embodiment, any one nucleotide of the control is in the hybridizing region of exactly two probes.

25 In another embodiment, a set of probes is complementary to regions of the control nucleic acid such that at least one nucleotide of the control separates each adjacent region, therefore a spacing of at least one nucleotide separates each adjacent probe in the set when hybridized. In this embodiment, at least a second set of probes similarly is complementary to regions of the control such that at least one nucleotide separates each adjacent region, each probe of the second being complementary to a region of the control containing the one or more nucleotides which separate adjacent probes of the first set. This will be referred to as

the nucleotide spacing embodiment. In a preferred embodiment, a single nucleotide separates each adjacent region in each probe set.

Any means of determining the melting temperature of a probe may be used. Several means of determining the melting temperature of hybridized nucleotides are known in the art.

In a preferred embodiment, fluorescent double-stranded DNA specific dyes are used to determine the melting temperature. In this embodiment, the fluorescent output of a sample of probe hybridized with target or control is monitored as the sample is heated. The fluorescent output is plotted against the temperature to determine the melting temperature of the probe from the target or control.

In another preferred embodiment, probes comprising a fluorescent label are used. For example, the hybridization of 5'-fluorescein-labeled probes, and melting thereof, can be observed when the fluorescein label is in close proximity to a guanosine residue on the complementary strand, due to the inherent quenching of fluoresceine signal by guanosine. (See, Crocket and Wittwer, *Anal. Biochem.*, in press (2001), incorporated herein in its entirety).

A difference in melting temperature of a probe from the target as compared with the melting temperature of an identical probe from the control (ΔT_m) indicates an alteration in the target sequence as compared to the control. A ΔT_m of a second probe localizes the alteration to the region of the target in which the two probes overlap. The difference between ΔT_m of two overlapping probes ($\Delta \Delta T_m$) indicates the exact position of the base change in the overlapping region because this number is unique for each residue location for each type of residue change, as further described below.

The magnitude of ΔT_m for a given probe also indicates the nature of a base change which is present. That is, the type of residue (e.g., adenine, guanosine, thymine, cytosine) that is different in the target as compared with the control may be determined. In a preferred embodiment, determination of the residue change in a target as compared to a control is made using a T_m profile map, as further described below. A " T_m profile map" is a table describing the anticipated ΔT_m for each possible residue change under consideration for each residue with which a probe hybridizes. The T_m profile map may be generated from experimental data for specific sequence alterations in a control sequence or from predicted T_m 's using thermodynamic calculations (See, Santa Luci et al., *Proc. Nat. Acad. Sci. USA* 95:1460-1465 (1998) and Schutz et al., *BioTechniques* 27:1218-1224 (1999)).

In a preferred embodiment, determination of the residue change in a target as compared to a control is made using ΔT_m of two or more probes. Example 3 further illustrates determination of the location and type of a single nucleotide change in a target as compared to a control.

In a preferred embodiment, the rate at which the temperature is increased during the determination of a melting temperature for a probe is the same for all probes and for each probe in determining the melting temperature from the target and from the control. In an alternate embodiment, the melting temperature for a probe of interest from a given target is determined at two or more rates of increasing temperature.

In a preferred embodiment, melting temperatures of a probe from one or more targets having known alterations are determined to produce a T_m -alteration profile or ΔT_m map for the probe.

A " T_m profile map" is a table describing the anticipated ΔT_m for each possible residue change under consideration for each residue with which a probe hybridizes. The T_m may be determined for alterations at different locations, as well as alterations involving different nucleotide substitutions. Additionally, the T_m for each of two or more different rates of increasing temperature may be determined for one or more alterations, to produce a T_m -alteration-rate profile. The T_m profile map may also be generated from predicted T_m 's using thermodynamic calculations, as is well known in the art. (See, Santa Luci et al., *Proc. Nat. Acad. Sci. USA* 95:1460-1465 (1998); Schutz et al., *BioTechniques* 27:1218-1224 (1999); and Peyret et al., *Biochemistry* 38(12):3468-3477 (1999), each of which is incorporated herein in its entirety).

The differences in ΔT_m between the two probes is used to determine the location within the overlap region of the two probes. Alternatively, the ΔT_m for each probe may be determined at two or more rates of increasing temperature.

In the single nucleotide spacing embodiment using two sets of probes, usually two probes will hybridize to the region having any single base mutation and each of said probes will have a ΔT_m . However, in the case where a single base change occurs in the gap between the hybridization region of two probes of one probe set, only one probe of the other probe set will have a ΔT_m . Otherwise, when the ΔT_m of a first probe is subtracted from the ΔT_m of a second overlapping probe, the difference in ΔT_m between the overlapping probes ($\Delta \Delta T_m$) will increase for each alteration that is closer to the end of the first probe. Alternatively, the location of the alteration may be determined from the ΔT_m -alteration-rate profile for either or both of the first and second probes.

In addition to the location of the alteration, the nature of a nucleotide substitution may be determined as described above. In one embodiment, a T_m -alteration profile for one or both overlapping probes with a ΔT_m is used to determine the substitution.

5 In one embodiment, the invention is implemented in a nucleotide solid support array. Such arrays are well known in the art. An example of such an array is described in U.S. Patent No. 5,571,639, incorporated herein in its entirety.

10 In a preferred embodiment, the determination of the T_m of the probes for a given target and control is done simultaneously.

15 In a preferred embodiment, the invention is automated. In this embodiment specialized hardware and software programs may be applied to control the automation of the invention. One system useful in the present invention is the commercially available LightCycler™, made by Idaho Technology (Wittwer et al., *BioTechniques* 22:176-181 (1997), incorporated herein in its entirety). This apparatus is capable of rapid thermal cycling necessary for amplification of nucleic acid for procedures such as PCR. This apparatus is also capable of providing a probe/control and/or probe/target sample with a prescribed temperature within a very broad range and change temperatures at a prescribed rate. The LightCycler™ is also capable of
20 measuring fluorescence in a sample and changes in fluorescence coincidentally with changing temperature for T_m determinations.

25 The present invention finds uses in many fields, including but not limited to genetics, immunology, infectious disease, oncology, epidemiology and forensics. Such uses include, but are not limited to, identifying mutations in tumorigenic material, identifying inheritable genetic disease and guiding treatment for such diseases. Additionally, the invention may be used for identifying allelic variants, identifying sources of biological samples and determining paternity.

30 It will be apparent to one of ordinary skill in the art that many other applications of the present invention are possible. All references cited herein are incorporated in their entirety.

The following examples are provided for illustrative purposes. It is understood that these examples in no way serve to limit the true scope of the invention.

35

EXAMPLES

Example 1

Asymmetric PCR is performed by rapid cycling techniques in a reaction volume of 10 μ l with dNTP's at 200 μ M each, one primer at 0.5 μ M, the other primer at 0.05 μ M, 3 mM Mg^{2+} , 50 mM Tris (pH 8.3), 500 mg/ml BSA. 0.4 U Taq polymerase, and 10 ng of genomic DNA. After 50 cycles of 94° C for 0 seconds, 55° C for 0 seconds and 72° C for 10 seconds, the sample is cooled to 40° C and a high resolution melting curve is performed by heating at 0.05° C per second in the presence of fluorescent double-stranded DNA-specific dye.

Melting curves are acquired for all probes and the first derivative of each curve is calculated and plotted as -dF/dT (the negative first derivative of fluorescence with respect to temperature). T_m 's are determined as the peak of each derivative curve by Gaussian fitting. ΔT_m 's (the difference in T_m between control and test samples) and $\Delta\Delta T_m$'s (the difference in ΔT_m when using different probes) are calculated and used to identify sequence alterations in the test sample as compared to the control sample.

Example 2

Single stranded nucleic acid of a control and a target are separately produced by asymmetric PCR, SDA or 3SR. After amplification with a biotinylated primer, the amplified control and target are purified by attachment to streptavidin-coated magnetic beads. The purified single strand nucleic acid is placed in probe wells or a probe chip and analyzed by melting as in Example 1.

Example 3

Single nucleotide polymorphism (SNP) screening and identification.

Two sets of 15-mer nucleic acid probes are produced for use in the nucleotide spacing embodiment described above, wherein the number of nucleotides separating adjacent complementary regions for each probe set is one (see Fig. 2). If a ΔT_m is observed for one probe then the SNP is localized to the specific base that is unique for that probe. The magnitude of ΔT_m identifies the specific base change (see Fig. 3). If a ΔT_m is observed for two probes, then the SNP is localized to the base positions where the overlap between probes occurs. The specific position and base change is identified from a T_m alteration profile map that is either measured experimentally or predicted from thermodynamic calculations

An example of a partial T_m alteration profile for the overlapping portion of 2 probes is shown in Table 1. There are seven possible positions that overlap between these 15-mer probes. The partial T_m alteration map of Table 1 shows all possibilities for SNP changes for an A in the control DNA (a complete map would contain 3 other Tables of equal size for C, G, and T, but the principle is the same). The values of $\Delta T_m(1)$, $\Delta T_m(2)$, and $\Delta \Delta T_m$ in the Table are used to identify the position and precise SNP that is present.

Table 1. Sequence Map for base A in the control sequence

Position								
	1	2	3	4	5	6	7	Mismatch
$\Delta T_m(1)$	5	5	5	5	4	3	1	
$\Delta T_m(2)$	1	3	4	5	5	5	5	A:G
$\Delta \Delta T_m$	4	2	1	0	-1	-2	-4	
$\Delta T_m(1)$	10	10	10	10	8	6	2	A:C
$\Delta T_m(2)$	2	6	8	10	10	10	10	
$\Delta \Delta T_m$	8	4	2	0	-2	-4	-8	
$\Delta T_m(1)$	8	8	8	8	7	4	2	A:T
$\Delta T_m(2)$	2	4	7	8	8	8	8	
$\Delta \Delta T_m$	6	4	1	0	-1	-4	-6	

With high-resolution determination of the T_m s, every possible SNP (there are 84 possibilities in the 7 positions, 21 are shown in Table 1) can be identified by a unique value of $\Delta T_m(1)$, $\Delta T_m(2)$, and $\Delta \Delta T_m$. Seldom is such precision required, and often the need is to distinguish between several sequence possibilities. In this case, a limited sequence map can be construction with only the sequence alterations that need to be screened. In this case, lower resolution equipment can be used to discriminate between the possibilities (not necessarily limited to SNPs). Another method to increase discrimination is to use different melting temperature rates (for example 0.1°C/sec and 0.5°C/sec to estimate T_m (Ririe et al., *Anal. Biochem.* 245:154-160 (1997), incorporated herein in its entirety). Alternately, both heating and cooling curves can be obtained to estimate T_m by melting and annealing (Gundry et al., *Genetic Testing* 3:365-370 (1999), incorporated herein in its entirety). When 2 different

apparent T_m 's are measured for each probe/target combination, the matrix will approximately double in size, providing twice as many parameters for distinguishing each possible genotype.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2